

Differential Expression of microRNA-2b with Potential Target Coding P25 in the Fifth Instar Larvae Posterior Silk Gland of the Silkworm¹

Yong Huang^{a, c}, Quan Zou^b, Xing Jia Shen^{a, c}, Shun Ming Tang^{a, c},
Sheng Peng Wang^{a, c}, and Qiao Ling Zhao^{a, c}

^a The Key Laboratory of Silkworm and Mulberry Genetic Improvement, Ministry of Agriculture,
Jiang Su University of Science and Technology, 212018, Jiangsu Province, PR China

^b School of Information Science and Technology of Xiamen University, 361005, Fujian Province, PR China

^c Sericultural Research Institute, Chinese Academy of Agricultural Sciences, 212018, Jiangsu Province, PR China;
e-mail: shenxj63@yahoo.com.cn

Received August 23, 2010; in final form, September 24, 2010

Abstract—MicroRNAs (miRNAs) are a class of non-protein coding small RNAs that regulate a gene expression at the post-transcriptional level. Using in silico screening, we found that the 3'-untranslated regions of the *P25* gene mRNA are perfectly complementary to nucleotides 2–8 at the 5' end of the miRNA-2b (miR-2b). The expression of miR-2b and the *P25* gene in posterior silk gland of the fifth instar larval silkworm was investigated using real-time PCR detection method. The results indicated that expression of the *P25* gene was very high in the posterior silk gland during the fifth instar larvae, whereas a level of miR-2b sharply decreased until reaching the lowest one on the 8th day. The expression patterns of miR-2b and *P25* gene indicate that miR-2b might act as a fine-tuning regulator of expression of the *P25* gene at the post-transcriptional level.

DOI: 10.1134/S0026893311040133

Keywords: microRNA, silkworm, in silico, *P25* gene.

INTRODUCTION

MicroRNAs (miRNAs) are small non-coding RNAs, which act as important post-transcriptional gene regulators and control the expression of target genes in eukaryotes. miRNAs are expressed as long hairpin-forming precursor RNAs that are cleaved into partially double-stranded RNAs, which are then further processed into mature miRNAs (22 nucleotides) [1–4]. Mature miRNAs recognize their target mRNAs by base-pairing interactions between nucleotides 2–8 of the miRNA (the seed region) and the complementary nucleotides in the 3'-untranslated region (3'-UTR) of the mRNAs. miRNAs inhibit gene expression by targeting mRNAs for translational repression or cleavage [5–7].


The silkworm (*Bombyx mori*) is one of the most economically important insects in many developing countries because of its large-scale cultivation for silk production. Silk fibroin produced by the silkworm consists of a heavy chain (F-H), a light chain (F-L), and *P25* (which is a glycoprotein of about 30 kDa) [8–10]. The expression of the three fibroin genes in silkworm is controlled at the transcription level during larval

developmental stages [11]. Silk fibroin has been reported to be specifically synthesized in the posterior silk gland (PSG) cells of silkworm and secreted into its lumen in a form of 2.3-MDa protein complex, in which the F-H, F-L, and *P25* are present in 6 : 6 : 1 molar ratios per elementary unit of fibroin [12]. However, molar ratios of corresponding mRNAs greatly differ from the protein composition in the PSG cell [13]. Based on the results, we assumed that expression of the three fibroin genes might be regulated at the post-transcriptional level.

Previous studies demonstrated that the expression of fibroin genes is mainly co-regulated by a series of *trans*-acting factors and the corresponding *cis*-acting elements at the transcriptional level [14]. Recently, it has been reported that at least 4 miRNAs (miR-33, miR-190, miR-276, and miR-7) might play an important role in the regulation of fibroin L-chain transcripts [15]. However, few studies have been reported on miRNAs involved in the regulation of the *P25* gene. Based on the fact mentioned above, we chose the *P25* gene as our research object. In silico search for potential miRNAs probably interacting with the 3'-UTR of the *In silico* mRNA showed that the miR-2b seed region is perfectly complementary to the target. Thus,

¹ The article is published in the original.

RNAhybrid prediction:



RNA22 prediction:




Fig. 1. In silico derived base pairing between miR-2b and the 3'-UTR mRNA of *P25* gene. The miR-2b and target *P25* are partitioned into two parts, the seed region and out-seed region. The seed sequence corresponds to seven nucleotides at positions 2–8 of the miRNA sequence. The seed site has perfect complementarity with the miRNA 5'-end that is indicated by black bold letters.

miR-2b may be involved in the post-transcriptional regulation of the expression of the *P25* gene. In order to explore the miRNA regulation mechanism at the post-transcriptional level, we compared the expression of miR-2b with that of *P25* gene in the posterior silk gland of the fifth instars larvae using real-time PCR assay. The results indicated that miR-2b was probably to be involved in the post-transcriptional regulation of the *P25* gene.

EXPERIMENTAL

Materials. A bivoltine strain (DaZao) of silkworm was used in the experiments. The posterior silk glands of the fifth instar larvae were separately dissected at 1 day, 3 day, 5 day, 7 day and 8 day, immediately frozen in liquid nitrogen, and then stored at -80°C .


Referenced sequence data. Mature silkworm miRNAs were downloaded from the miRNA database (<http://microrna.sanger.ac.uk>; Release 14.0, September 2009). The 3'-UTR of *P25* gene mRNA was downloaded from NCBI (<http://www.ncbi.nlm.nih.gov/>

[sites/entrez?db=unigene](http://www.ncbi.nlm.nih.gov/sites/entrez?db=unigene)). The potential target sites were detected using RNAhybrid and RNA22 tool. At the 5'-end of the miRNA there is a region called “seed”, which is centered on nucleotides 2–8 [16, 17]. miRNA targets were difficult to predict since miRNA : mRNA duplexes often contain several mismatches, gaps and G:U or T:U base pairs in many positions according to different target prediction methods [18]. RNAhybird is an extension of classical RNA secondary structure prediction method [19]. It calculates the energetically most favorable hybridization of a miRNA and its target mRNA rapidly and accurately, which also allowing the user to specify a portion of the miRNA that should form a perfect helix, corresponding to the seed site. RNA22 is distinct from previously reported methods in that it doesn't consider cross-species sequence conservation. It is a pattern-based method for miRNA target prediction [20]. It makes use of miRNA patterns to identify “target islands”, regions of the mRNA sequence that have a higher likelihood to contain binding sites for miRNAs.

Sequences of stem-loop RT primers, forward primers, and reverse primers for real-time PCR

Gene Name	Primer	Sequence
miRNA-2b	RT	5'-GCGTGGTCCCGACCACCACAGCCGCCACGACCACGCACTCAA-3'
	Forward	5'-GCGTATCACAGCCAGCTTTG-3'
	Reverse	5'-TCCCGACCACCACAGCC-3'
5S	RT	5'-CTCAACTGGTGTCGTGGAGTCGGCAATTCAGTTGAGAAGCCTACGCTCTGGAATACCGGGTGC-3'
	Forward	5'-GCTCTGGAATACCGGGTGC-3'
	Reverse	5'-TCCCGACCACCACAGCC-3'
<i>P25</i>	Forward	5'-CCCTGCTACTTGGACGATT-3'
	Reverse	5'-GATTATGGTCGACGTAGGTG-3'

RNAhybrid prediction:



RNA22 prediction:




Fig. 2. In silico derived base-pairing between miRNAs and the *P25* 3'-UTR. Black bold letters indicates the 'seed region' of the binding site.

Total RNA isolation. Total RNA was isolated from posterior silk gland tissues of silkworm fifth instar larvae (see Materials) using Trizol (Invitrogen) according

to the manufacturer's instructions. Contaminating genomic DNA was removed by DNase I (Takara, Dalian). The purity of extracted RNA was determined by a UV spectrophotometer NanoDrop™ 1000 (Thermo Scientific). Ratios of UV 260/280 were between 1.8 and 2.1 for all of analyzed RNA samples. The total RNA concentration was determined by measuring the absorbance at 260 nm.

Real-time PCR quantification. cDNA was synthesized from a total of 500 ng of RNA in a 20-μl reaction system by using specific stem-loop primers for miRNA obtained from a commercial service (Takara) and a common prime script™ RT reagent kit (Takara) for the *P25* gene. The cDNAs were diluted 10 times. Real-time PCR was performed in 20 μl mixture containing 1 μl of cDNA, 800 nM forward and reverse primers, 10 μl of the 2× SYBR green PCR master mix (Takara) [21]. Real-time PCR primers used in this experiment are indicated in the table. The reaction was performed in triplicate on an opticon lightcycler (Bio-Rad Laboratories). The reactions were amplified for 15 s at 95°C and 30 s at 57°C for 40 cycles.

Statistical analysis. Melting curves for each PCR were carefully monitored to avoid nonspecific amplification. Expression levels of the target genes were normalized against the expression level of the 5S rRNA gene by Option monitor analysis software (MJ Research). Default threshold settings were used to determine the threshold cycle data. The relative amount of each miRNA to 5S rRNA was described by using the equation $2^{-\Delta Ct}$, where $\Delta Ct = Ct(\text{miRNA}) - Ct(5S \text{ rRNA})$ [22, 23]. miRNA expression data were normalized by the levels of 5S rRNA. Data were presented as the mean value $\pm$ SD from two or three separate experiments. The *p*-values lower than 0.05 were considered as significant.

RESULTS

In silico Prediction of the *P25* mRNA 3'-UTR's Base-Pairing with the miRNA 5' Seed Region

Although animal miRNAs tend to show imperfect base-pairing in the 3'-UTRs of their target transcripts, a 7- or 2–8-nucleotide seed sequence starting from the second nucleotide at the 5'-end of miRNAs with their target mRNA that leads to decreased transcription and/or translation repression [24, 25]. Here, two target prediction tools, the RNAhybrid and RNA22, were used to predict miRNAs through the silkworm miRNA database and to find the seed sites complementary to the 3'-UTR of *P25* gene mRNA (Fig. 1). The results manifested that miR-2b meets the In addition, through searching the new miRNA database (Release 15, April 2010), four miRNAs potentially directed to the 3'-UTR of *P25* gene mRNA were found (Fig. 2). These miRNAs are complimentary to

different regions within the 3'-UTR and therefore might affect the expression of *P25* gene.

Expression of miR-2b and *P25* Gene

An expression level of miR-2b varied across the development cycle of the posterior silk gland in the fifth instar larvae. It was the highest at the first day and then sharply decreased during most of the 8 development days (Fig. 3a).

Fibroin genes were expressed exclusively highly in the posterior silk gland of the fifth instar larvae [26, 27]. Similarly, real-time PCR analysis revealed a high level of *P25* gene expression in the posterior silk glands of the fifth instar larvae (Fig. 3b). Its expression was the lowest on the first day; obviously increased at day 3; reached maximal value on the 5 day, and then decreased on the 7 and 8 days. On the base of results obtained we hypothesize that expression of the *P25* gene might be regulated at the transcription level during specific developmental stages in the posterior silk gland of the fifth instar larvae. Thus, we assumed that expression of the *P25* gene might be regulated at the transcription level during specific developmental stages in the posterior silk gland of the fifth instar larvae.

DISCUSSION

At present, there are certain characteristics of the interactions between miRNAs and their targets which current algorithms generally use: (1) the complementarity between miRNAs and their target sites exists; (2) the 5'-end of a miRNA binds to the target mRNA stronger than the 3'-end does. In addition to these basic principles, different prediction methods impose different restrictions on the algorithms to achieve optimization in accordance with their respective rules [28–31]. In order to identify the miRNA targets in the 3'-UTR of *P25* gene mRNA for evaluating miRNA-target association, two computational target-prediction approaches were used. The results revealed that the 2–8 nucleotides of miR-2b seed region were full complementary to the identically located sequence within *P25* mRNA 3'-UTR (Fig. 1). The outcome strongly indicates that miR-2b might be involved in the post-transcriptional regulation of the *P25* gene expression. Meanwhile, among the new identified silkworm miRNAs there are more possible miRNAs which may regulate the expression of the *P25* gene (Fig. 2). These prediction results are required further test and verification in vivo and in vitro experiments using methods of miRNA over-expression and knocked-down.

It was shown that some miRNAs could up-regulate translation besides their down-regulating role [32–34]. In this paper, we used real-time RT-PCR method




Fig. 3. Expression of miR-2b (a) and *P25* gene (b) in the posterior silk gland of the fifth instar larvae. Data are represented as the mean value $\pm$ SD from three independent experiments.

for evaluating the *P25* gene and miR-2b expression. As one can see from the data presented in Figs. 3a and 3b, expression of *P25* gene increased significantly from day 1 to day 3, and reached the highest value at the 5 day, whereas the expression of miR-2b, on the contrary, decreased at that time interval. Based on the results we suppose that there might be a certain regulation relationship of expression between miR-2b and the *P25* gene. Obviously, our hypothesis will further need to be verified in vivo experiments. It is known that miRNAs are often expressed in a strict tissue-specific manner during development stages [35–37]. As one can see in Fig. 3b, the expression of *P25* gene displayed obviously the down-regulation from day 7 probably because a lot of fibroin genes would be activated and translated into fibroin proteins in this development stage.

To sum up, through combination with two methodologies, the computational prediction for a search of *P25* 3'-UTR-targeted miRNAs and real-time RT-PCR for testifying model obtained, we conclude that miR-2b might play a biological role in regulating the *P25* gene expression.

ACKNOWLEDGMENTS

This work was partly supported by the National Natural Science Foundation of China (NSFC31072083); Innovation Foundation for Graduate Students of Jiangsu Province and the Natural Science Foundation of China (61001013).

REFERENCES

- Ambros V. 2004. The functions of animal microRNAs. *Nature*. **431**, 350–355.
- Lee R.C., Feinbaum R.L., Ambros V. 1993. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell*. **75**, 843–854.
- Brennecke J., Stark A., Russell R.B., Cohen S.M. 2005. Principles of microRNA-target recognition. *PLoS Biol.* **3**, e85.
- Huang Y., Zou Q., Wang S.P., Tang S.M., Zhang G.Z., Shen X.J. 2010. The discovery approaches and detection methods of microRNAs. *Mol. Biol. Rep.* doi: 10.1007/s11033-010-0532-1.
- He L., Hannon G.J. 2004. MicroRNAs: small RNAs with a big role in gene regulation. *Nature Rev. Genet.* **5**, 522–531.
- Lewis B.P., Burge C.B., Bartel D.P. 2005. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*. **120**, 15–20.
- Huang Y., Shen X.J., Zou Q., Wang S.P., Tang S.M., Zhang G.Z. 2010. Biological functions of microRNAs: A review. *J. Physiol. Biochem.* doi: 10.1007/s11033-010-0532-1.
- Couble P., Moine A., Garel A., Prudhomme J.C. 1983. Developmental variations of a nonfibroin mRNA of *Bombyx mori* silk gland, encoding for a low-molecular-weight silk protein. *Dev. Biol.* **97**, 398–407.
- Sprague K.U. 1975. The *Bombyx mori* silk proteins: characterization of large polypeptides. *Biochemistry*. **14**, 925–931.
- Yamaguchi K., Kikuchi Y., Takagi T., Kikuchi A., Oyama F. 1989. Primary structure of the silk fibroin light chain determined by cDNA sequencing and peptide analysis. *J. Mol. Biol.* **210**, 127–139.
- Kimura K., Oyama F., Ueda H., Mizuno S., Shimura K. 1985. Molecular cloning of the fibroin light chain complementary DNA and its use in the study of the expression of the light chain gene in the posterior silk gland of *Bombyx mori*. *Experientia*. **41**, 1167–1171.
- Inoue S., Tanaka K., Arisaka F., Kimura S., Ohtomo K. 2000. Silk fibroin of *Bombyx mori* is secreted, assembling a high molecular mass elementary unit consisting of H-chain, L-chain, and P25, with a 6 : 6 : 1 molar ratio. *J. Biol. Chem.* **275**, 40517–40528.
- Grzelak K. 1995. Control of expression of silk protein genes. *Comp. Biochem. Physiol. B: Biochem. Mol. Biol.* **110**, 671–681.
- Okamoto H., Ishikawa E., Suzuki Y. 1982. Structural analysis of sericin genes: Homologies with fibroin gene in the 5' flanking nucleotide sequences. *J. Biol. Chem.* **257**, 15192–15199.
- Cao J., Tong C., Wu X., Lv J., Yang Z. 2008. Identification of conserved microRNAs in *Bombyx mori* (silkworm) and regulation of fibroin L chain production by microRNAs in heterologous system. *Insect. Biochem. Mol. Biol.* **38**, 1066–1071.
- Huang Y., Zou Q., Song H., Song F., Wang L., Zhang G., Shen X. 2010. A study of miRNAs targets prediction and experimental validation. *Protein Cell*. **1**, 979–986.
- Min H., Yoon S. 2010. Got target? Computational methods for microRNA target prediction and their extension. *Exp. Mol. Med.* **42**, 233–244.
- Mendes N.D., Freitas A.T., Sagot M.F. 2009. Current tools for the identification of miRNA genes and their targets. *Nucleic Acids Res.* **37**, 2419–2433.
- Rehmsmeier M., Steffen P., Hochsmann M., Giegerich R. 2004. Fast and effective prediction of microRNA/target duplexes. *RNA*. **10**, 1507–1517.
- Miranda K.C., Huynh T., Tay Y., Ang Y.S., Tam W.L. 2006. A pattern-based method for the identification of microRNA binding sites and their corresponding heteroduplexes. *Cell*. **126**, 1203–1217.
- Yu X., Zhou Q., Li S.C., Luo Q., Cai Y. 2008. The silkworm (*Bombyx mori*) microRNAs and their expressions in multiple developmental stages. *PLoS One*. **3**, e2997.
- Schmittgen T.D., Livak K.J. 2008. Analyzing real-time PCR data by the comparative C(T) method. *Nature Protoc.* **3**, 1101–1108.
- Mardanov E.S., Zamchuk L.A., Ravin N.V. 2007. The 5' untranslated region of the maize alcohol dehydrogenase gene provides efficient translation of mRNA in plants under stress conditions. *Mol. Biol. (Moscow)*. **41**, 914–919.
- Lai E.C., Tomancak P., Williams R.W., Rubin G.M. 2003. Computational identification of *Drosophila* microRNA genes. *Genome Biol.* **4**, R42.
- Lewis B.P., Shih I.H., Jones-Rhoades M.W., Bartel D.P., Burge C.B. 2003. Prediction of mammalian microRNA targets. *Cell*. **115**, 787–798.
- Datta A., Ghosh A.K., Kundu S.C. 2001. Differential expression of the fibroin gene in developmental stages of silkworm, *Antheraea mylitta* (Saturniidae). *Comp. Biochem. Physiol. B: Biochem. Mol. Biol.* **129**, 197–204.
- Liu S., Zhang L., Li Q., Zhao P., Duan J., Cheng D., Xiang Z., Xia Q. 2009. MicroRNA expression profiling during the life cycle of the silkworm (*Bombyx mori*). *BMC Genomics*. **10**, 455.
- Chaudhuri K., Chatterjee R. 2007. MicroRNA detection and target prediction: Integration of computational and experimental approaches. *DNA Cell Biol.* **26**, 321–337.
- Li L., Xu J., Yang D., Tan X., Wang H. 2010. Computational approaches for microRNA studies: A review. *Mamm. Genome*. **21**, 1–12.
- Sethupathy P., Megraw M., Hatzigeorgiou A.G. 2006. A guide through present computational approaches for the identification of mammalian microRNA targets. *Nature Methods*. **3**, 881–886.

31. Xia W., Cao G., Shao N. 2009. Progress in miRNA target prediction and identification. *Sci. China C Life Sci.* **52**, 1123–1130.
32. Vasudevan S., Tong Y., Steitz J.A. 2007. Switching from repression to activation: microRNAs can up-regulate translation. *Science*. **318**, 1931–1934.
33. Vasudevan S., Steitz J.A. 2007. AU-rich-element-mediated upregulation of translation by FXR1 and argonaute 2. *Cell*. **128**, 1105–1118.
34. Buchan J.R., Parker R. 2007. The two faces of miRNA. *Science*. **318**, 1877–1878.
35. Ason B., Darnell D.K., Wittbrodt B., Berezikov E., Kloosterman W.P. 2006. Differences in vertebrate microRNA expression. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 14385–14389.
36. Aboobaker A.A., Tomancak P., Patel N., Rubin G.M., Lai E.C. 2005. *Drosophila* microRNAs exhibit diverse spatial expression patterns during embryonic development. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 18017–18022.
37. Lagos-Quintana M., Rauhut R., Yalcin A., Meyer J., Lendeckel W. 2002. Identification of tissue-specific microRNAs from mouse. *Curr. Biol.* **12**, 735–739.